

GAPDH Polyclonal Antibody

Catalogue No*:	AB0049-200	Price/Unit*:	320.00
Source:	3	Quantity/unit size*:	400 g
Description:	Goat polyclonal to GAPDH (glyceraldehyde 3-phosphate dehydrogenase). GAPDH catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The enzyme exists as a tetramer of identical chains. Alternative names: glyceraldehyde 3-phosphate dehydrogenase, glyceraldehyde-3-phosphate dehydrogenase, G3PD, GAPD, HGNC:4141, GAPDH antibody.		
Immunogen:	Purified recombinant peptide derived from within residues 240 aa to the C-terminus of human GAPDH produced in <i>E. coli</i> .		
Specificity:	Detects a band of 37 kDa by Western blot in the following human (293A, HMEC-1, U-118, HaCat), rat (TR-iBRB), mouse (AtT-20, Hepa), canine (D17) and monkey (COS-7) whole cell lysates.		
Reactivity:	Reacts against human, rat, mouse, zebrafish, canine and monkey proteins.		
Usage:	Western blot 1:500-1:5,000 Immunofluorescence 1:50-1:250 Immunohistochemistry (paraffin) 1:200-1:1,000 Immunohistochemistry (frozen) 1:200-1:1,000		

Reactivity Chart



Sample	Western Blot	Immuno-fluorescence	Immuno-Histochemistry (p)	Immuno-Histochemistry (f)
human	+++	+++	+++	+++
rat	+++	+++	+++	+++
mouse	+++	+++	+++	+++
zebrafish	+++	ND	ND	ND
canine	+++	+++	+++	+++
monkey	+++	+++	+++	+++

Storage:	Store at -20 C for long term storage. Store at 2-8 C for up to one month.
Special instructions:	Avoid freeze/thaw cycles.

References:

1. Martins SGR, MSc Thesis, NOVA University of Lisbon, Portugal 2019
2. Ferreira JV, Rosa Soares A, Ramalho JS, et al. *PLoS One* 2019 Oct. PMID:31613922
3. Alenquer M, Vale-Costa S, Etibor TA, et al. *Nat Commun* 2019 Apr. PMID:30967547
4. Sanz P, Evans RD, Briggs DA, et al. *J Cell Sci* 2019 Apr. PMID:30898842
5. Aires ID, Boia R, Rodrigues-Neves AC, et al. *Glia* 2019 Jan. PMID:3066709
6. Barbeitos JP, MSc Thesis, University of Coimbra, Portugal 2018
7. Alenquer M, Vale-Costa S, Sousa AL, et al. *bioRxiv* 410373; Sept 2018
8. Alzahofi N, Robinson CL, Welz T, et al. *bioRxiv* 314153; May 2018
9. Ribeiro ST, Tesio M, Ribot JC, et al. *Leukemia* 2017 Jul. PMID:27899804
10. Santarino IB, Viegas MS, Domingues NS, et al. *Sci Rep* 2017 Jul. PMID:28724916
11. Ribeiro STF, PhD Thesis, University of Lisbon, Portugal 2017
12. Robinson CL, Evans RD, Briggs DA, et al. *J Cell Sci* 2017 May. PMID:28490438
13. Ramalho AR, Toscano A, Pereira P. et al. *Rev Port Cardiol* 2017 May. PMID:28479269
14. Vale-Costa S, Alenquer M, Sousa AL, et al. *J Cell Sci* 2016 Mar. PMID:26940915
15. Encarnao M, Espada L, Escrevente C, et al. *J Cell Biol* 2016 Jun. PMID: 27325790
16. Ferreira JV, Soares AR, Ramalho JS, et al. *Sci Rep* May 2015. PMID:25958982
17. Ferreira RRS, MSc Thesis, University of Coimbra, Portugal 2015
18. Paiva RA, MSc Thesis, University of Coimbra, Portugal 2015
19. Casalou C, Seixas C, Portelinha A, et al. *J Cell Sci* 2014 Jun. PMID:24777479
20. Ribeiro-Rodrigues TM, Catarino S, Marques C, et al. *FASEB J* 2014 Nov. PMID:25070368
21. Moreiras HAF, MSc Thesis, University of Lisbon, Portugal 2014

For research use only, not for diagnostic use



Anti-GAPDH Ab at 1/2,500 dilution; lysates at 50 µg per lane; Rabbit polyclonal to goat IgG (HRP) at 1/10,000 dilution;

Immunofluorescence – anti-GAPDH Ab in Hepa1-6 cells at 1/50 dilution; cells were fixed with methanol;